

High p_T in Nuclear Collisions at the SPS, RHIC, and LHC

P. Jacobs ^a and M. van Leeuwen ^a

^aLawrence Berkeley National Laboratory,
1 Cyclotron Road,
Berkeley, CA
USA 94720

We review recent progress in the study of medium-induced modification of jet fragmentation in high energy nuclear collisions at the SPS and RHIC and present an outlook on jet physics at the LHC.

1. Introduction

Partonic energy loss is a potentially sensitive tomographic probe of matter produced in high energy nuclear collisions, generating observable modification of the fragmentation patterns of jets (“jet quenching”). Measurements at RHIC have revealed large medium-induced suppression at high transverse momentum (high p_T) of both the inclusive hadron yields [1,2] and of back-to-back hadron pairs [3]. The principal energy loss mechanism underlying these effects is commonly thought to be medium-induced gluon bremsstrahlung, which is expected to dominate collisional (elastic) energy loss for very energetic partons [4].

The effects of medium-induced radiation have been calculated in various frameworks: multiple soft scattering (BDMPS [5]), few hard scatterings (GLV [6]), and twist expansion (Wang and Guo [7]). In the case of multiple soft scattering the medium is characterized by a single transport coefficient $\hat{q} = \mu^2/\lambda$, where μ is the average momentum kick of a gluon interacting in the medium and λ is its mean free path. The gluon radiation spectrum is suppressed relative to the Bethe-Heitler spectrum due to coherence effects, leading to medium-induced radiated energy $\Delta E_{medium} \sim \alpha_s \hat{q} L^2$ [8,9]. Longitudinal expansion of the medium reduces the length dependence to $\Delta E_{medium} \sim L$ while finite partonic energy truncates the medium-induced radiation spectrum, resulting in an energy-dependent energy loss. Although the induced radiation spectrum differs in detail between the BDMPS and GLV approaches, the total energy loss is similar for comparable medium properties [9]. For conditions relevant to RHIC collisions the energy is lost in both cases to a moderate number (~ 3) of radiated gluons having moderate energy ($\sim 0.1 - 1$ GeV) [9].

This picture is conceptually appealing and pQCD-based calculations incorporating medium-induced bremsstrahlung reproduce much of the published data on high p_T hadron production in nuclear collisions. Nevertheless, it is important to ask to what extent the data *require* this description to be the correct one. Are its detailed predictions, such as the expected L^2 dependence of ΔE , observed? Alternatively, does *collisional* energy loss




Figure 1. Nuclear modification factor $R_{CP}(p_T)$ measured at the SPS and compared to model calculations [13]. Left panel: K_s^0 from NA57 [15]; Right panel: charged pions and protons from NA49 [16].

play a significant role in the finite kinematic regime of RHIC [10]?

We review recent measurements of partonic energy loss in hot matter, emphasizing results shown at this conference, and compare their systematic behavior to theoretical expectations. The variables at our disposal are energy, centrality and p_T dependence of high- p_T hadron production and correlations. We concentrate on the highest available p_T at mid-rapidity.

2. SPS

A long-standing puzzle concerns the magnitude of partonic energy loss in nuclear collisions at the SPS. WA98 data [11] indicated a large enhancement of inclusive π^0 production at $p_T \sim 3 - 4$ GeV/c in central Pb+Pb relative to p+p collisions when normalized per binary collision ($R_{AA}(p_T)$), although a suppression was observed in central relative to peripheral nuclear collisions ($R_{CP}(p_T)$, also normalized per binary collision) by the same experiment.

The p+p spectrum used as a reference for $R_{AA}(p_T)$ at the SPS was not directly measured but was based on an extrapolation from measurements at higher \sqrt{s} . Some of the datasets used in this extrapolation are only marginally consistent, leading to large uncertainties in the p+p reference spectrum. A recent re-evaluation of the extrapolation by d’Enterria [14] found a reduced $R_{AA}(p_T)$, more consistent with $R_{CP}(p_T)$. The medium density inferred from the new $R_{AA}(p_T)$ values is now also consistent with expectations from Bjorken energy density estimates [12,13].

New measurements of high- p_T hadron suppression at the SPS were presented at this conference. Fig. 1 shows $R_{CP}(p_T)$ for K_s^0 from NA57 [15] (left panel) and charged pions and protons from NA49 [16] (right panel) compared to radiative energy loss calculations. A marked Cronin effect is expected (“no E loss” on left panel) due to the steep p_T




Figure 2. Inclusive high p_T hadron suppression at RHIC. Left: $R_{AA}(p_T)$ for π^0 and charged hadrons in central Au+Au collisions from PHENIX [24]. The curves show suppression calculated in the single hard scattering approach [12]. Right: N_{part} dependence of $R_{AA}(p_T)$ at $p_T > 6$ GeV for charged hadrons from Cu+Cu and Au+Au collisions measured by STAR [25], compared to calculations in the multiple soft scattering approach [26].

spectrum. Introduction of energy loss in a medium, with gluon density scaling as $dN_{ch}/d\eta$ [13], results in good agreement between calculation and data.

These new data solve the high- p_T suppression puzzle at the SPS: the medium densities inferred from bulk multiplicity and high- p_T inclusive hadron measurements are consistent. However, significant theoretical uncertainties remain due to a potentially large Cronin effect at these lower energies.

3. RHIC: inclusive yields

Rich and initially unexpected phenomenology has emerged in the “intermediate p_T ” region ($p_T \sim 2 - 5$ GeV/c) of nuclear collisions at RHIC. The enhancement of baryon relative to meson yields [17,18], the approximate scaling of elliptic flow v_2 with the number of constituent quarks [19], and correlation measurements in this region [20] suggest an interplay between hadronization of hard partons and the bulk medium, as discussed extensively elsewhere at the conference [19,21]. Here we will concentrate on the region $p_T \gtrsim 6$ GeV, i.e. above the region of the anomalous enhancement of the baryon/meson ratio [17,18], to avoid these complex hadronization phenomena that may obscure the modification of fragmentation due to partonic energy loss.

Fig. 2, left panel, shows the most recent measurement of π^0 $R_{AA}(p_T)$ from the high-statistics 2004 Au+Au run at RHIC [24]. The kinematic reach now extends to $p_T \sim 20$ GeV/c, well beyond the intermediate p_T region. The observed p_T -independence of the suppression over this broad range is well described by partonic energy loss models in both the few hard scattering [12] (dashed and solid line) and multiple soft scattering [22,23] (not shown) limits, requiring medium densities $dN_g/dy \sim 1000 - 1200$ and $\hat{q} \sim 5 - 15$ GeV²/fm respectively. The transport coefficient \hat{q} and gluon density dN_g/dy in these models are time-averaged quantities. The connection between these averaged quantities

and the gluon or energy density of the medium at early time is discussed elsewhere [22,23].

The path length dependence of hadron suppression is a key test of its underlying mechanism. During the 2005 RHIC run, large samples of Cu+Cu collisions at $\sqrt{s_{\text{NN}}}=200$ GeV were collected to measure high- p_T particle production in this smaller system with similar statistical reach as the long 2004 Au+Au run. First analyses of a subset of the total Cu+Cu dataset were presented at this conference. Fig 2, right panel, shows the centrality dependence of $R_{AA}(p_T)$ for $p_T > 6$ GeV charged hadrons, from Cu+Cu collisions (squares), compared to existing results from Au+Au collisions (circles) [25]. The most significant comparison is to central Cu+Cu, where the Glauber calculation uncertainties are smaller than for the semi-peripheral Au+Au collision data with similar $N_{part} \sim 100$. The data indicate that the inclusive hadron suppression is a function solely of N_{part} , with no observable sensitivity to the shape of the reaction zone. The lines indicate phenomenological fits to characterize the N_{part} dependence of $R_{AA}(p_T)$. The data prefer a decrease with $N_{part}^{1/3}$, although the more commonly expected $N_{part}^{2/3}$ scaling is not strongly excluded. In general, the observed scaling behavior results from the combined effects of the spectrum shape, the collision geometry, and the path-length dependent energy-loss distribution. The gray bands indicate the results of a full calculation incorporating these effects [26], which reproduces the common suppression in Cu+Cu and Au+Au at the same N_{part} but gives slightly larger suppression at low N_{part} than observed in the data.

4. RHIC: correlations

Sensitivity of inclusive hadron suppression measurements to the properties of the medium is limited due to a surface bias: for a dense system, the observed hadrons are preferentially the fragments of partons produced near the surface and headed outwards, which suffer less than average energy loss[27,22]. The observed $R_{AA}(p_T) \sim 0.2$ in central Au+Au collisions is reproduced by calculations with a broad range of $\hat{q} \sim 5 - 15$ GeV²/fm [22], suggesting that its value is essentially geometric in origin. A more detailed view of the medium is obtained from back-to-back correlations of hadron pairs, where the surface bias can be reduced by increasing the p_T -threshold for associated particles on the away-side. This leads to counterbalancing biases from the requirements on the trigger and recoil, providing a new and more sensitive probe of the medium.

It has been known for some time that the back-to-back high p_T di-hadron yield is strongly suppressed in the most central Au+Au collisions [3]. However, the kinematic cuts of this first measurement ($p_T^{trig} > 4$ GeV/c for the trigger and $p_T^{assoc} > 2$ GeV/c for the associated hadrons) were relatively low, giving rise to a large combinatorial background. Due to the large background and the strong suppression, a differential measurement of the recoil yield was not possible and only an upper limit could be established. Lowering p_T^{assoc} reveals an excess and broadening of the recoil yield [28], to the extent that the correlation structure is compatible with expectations from simple momentum conservation without additional dynamic correlations. For these lower p_T^{assoc} , however, the backgrounds are yet larger and the uncertainties in the background yield and the flow modulation of the background make a quantitative measurement of the excess difficult. Qualitatively, these observations are compatible with a picture in which strong partonic energy loss in the core of the reaction volume softens the fragmentation. The response of the medium to the




Figure 3. Per-trigger correlated yield for d+Au and Au+Au collisions in two different centrality ranges, with $p_T^{trig} > 8$ GeV. The different rows show the correlation for different ranges in p_T of the associated particle [31].

energy loss then becomes a key issue, which was discussed extensively elsewhere at this conference [20,29,30].

New analyses from the long 200 GeV Au+Au run at RHIC now provide much greater p_T reach for correlation studies, for the first time extending beyond the intermediate p_T region. Fig. 3 shows azimuthal correlations of charged hadrons for $p_T^{trig} > 8$ GeV/c and varying threshold on p_T^{assoc} [31]. It is seen from the figure that the combinatorial background is negligible for the highest cuts and a clear back-to-back correlation signal emerges, for the first time enabling a quantitative *differential* measurement of partonic energy loss.

This is further explored in Fig. 4 [31], which shows distributions of the near- (left panel) and away-side (right panel) associated hadrons as a function of di-hadron fragmentation variable $z_T = p_T^{assoc}/p_T^{trig}$ [32], for $8 < p_T^{trig} < 15$ GeV trigger hadrons in d+Au and semi-central and central Au+Au collisions. The lines in the right-hand panel indicate an exponential fit to the d+Au data (solid line) which is scaled by 0.54 and 0.25 to match the semi-central and central Au+Au data respectively (dashed lines).

On the near side, no significant variation of the yield or the fragmentation distribution is seen between the systems. Calculations in Ref. [33] predict a strong enhancement of the associated yield in central collisions due to large energy loss and corresponding trigger bias effects, which is not observed in the data.

In contrast to the centrality invariance of the near-side correlation, a strong suppression of the away-side yield is found in central collisions at a level ($\sim 25\%$) that is numerically similar to $R_{AA}(p_T)$. This away-side suppression is however not accompanied by observable modification to either the longitudinal fragmentation distribution (lines in Fig. 4) or the




Figure 4. Di-hadron fragmentation distributions on the near- (left panel) and away-side (right panel) for $8 < p_T^{trig} < 15$ GeV [31] in d+Au collisions and Au+Au collisions at two different centralities. The solid line shows an exponential fit to the d+Au data. The dashed lines have the same slope, but are scaled down to match the Au+Au data.

azimuthal distribution (see Fig. 3 and [31]).

Suppression that is independent of z_T for $z_T > 0.4$ is in qualitative agreement with calculations by Wang [32], though quantitatively the measured suppression is stronger than predicted. Strong suppression could be accompanied by broadening of the recoil distribution in the event-averaged two particle correlation function, due either to large-angle emission of fragments from radiated gluons or to medium-induced acoplanarity of the dijets. In the multiple soft scattering approach the relation between energy loss dE/dx and induced acoplanarity has the general form [34]

$$-\frac{dE}{dx} = \frac{\alpha_s N_c}{8} \cdot \langle p_T^2 \rangle_{jet}, \quad (1)$$

where $\langle p_T^2 \rangle_{jet}$ refers to the momentum transverse to the initial direction of the hard parton that is acquired from interactions in the medium. Detailed calculations of the broadening using this approach are not yet available. A calculation for large energy loss in the GLV framework [35] predicts that induced radiation will dominate the recoil hadron distribution up to high $p_T \sim 10$ GeV/c, leading to strong azimuthal broadening. In contrast, Fig. 3 shows no significant angular broadening in this kinematic range (see also [31]).

Current model calculations invoke independent gluon emission to calculate integrated energy loss [23] and may not incorporate both radiative and elastic energy loss in a consistent way. An improved theoretical framework is likely required to calculate accurately the induced acoplanarity. On the experimental side, higher-order multi-particle correlations [20,29] may be able to distinguish broadening due to additional radiation and

induced acoplanarity of the jets, which is not possible with event-averaged two-particle correlations.

Significant suppression of the recoil yield without measurable changes in the longitudinal and azimuthal correlations would arise if this measurement is sensitive primarily to hadronic fragments of partons that had little or no interaction with the medium. This bias arises naturally in radiative energy loss calculations, which predict finite probability to emit *zero* gluons in a medium of finite length [9], with emission of a moderate energy gluon leading to sizeable suppression in the relatively high p_T region of this analysis. A quantitative evaluation of these effects, including a realistic nuclear geometry and path length dependent energy loss distributions, is given in Ref. [26], mainly in the context of single-hadron suppression. These calculations indicate that the probability for both the trigger and recoil jet to have little interaction with the medium is small, and a contribution is expected from partons that experience significant interaction with the medium. Refinement of these and other calculations within the constraints of the new data presented at this conference will help elucidate the mechanisms underlying jet quenching and the properties of the medium. A tantalizing prospect from these jet quenching studies is an experimentally measured upper bound on \hat{q} , which could constrain the number of underlying degrees of freedom of the medium [37].

5. Jet Physics at the LHC

The CERN LHC is currently scheduled to commission p+p collisions in 2007, with the first Pb+Pb run at $\sqrt{s_{NN}}=5.5$ TeV in 2008 [38]. The factor 30 increase in collision energy relative to RHIC generates a huge increase in kinematic and statistical reach for hard probes. Figure 5 shows the yield for various observables relevant to jet quenching studies [39,45] expected from one LHC year of Pb+Pb running (10^6 seconds) at nominal luminosity. Simple binary collision scaling ($\propto A^2$) from calculated p+p cross-sections has been applied, with no nuclear effects taken into account.

There will be statistically robust yields for jets well above $E_T \sim 200$ GeV, providing logarithmically large energy variation over which to study jet quenching effects. The huge statistics will enable the study of rare, perturbatively calculable fragmentation channels such as very hard hadron pairs with small angular separation, whose distributions may be modified by medium effects. The large yield of high energy jets raises the possibility that multi-particle or calorimetric (quasi-)full jet reconstruction can be used even in the presence of large backgrounds in central nuclear collisions. This potentially recovers the energy radiated in gluons, allowing relatively unbiased reconstruction of the jet energy and enabling complete characterization of jet quenching without the complications of strong trigger and geometric biases that are present in leading hadron and di-hadron measurements as performed at RHIC.

Jet reconstruction with good energy resolution is not straightforward in central nuclear collisions even at the LHC, however. For jets with $E_T \sim 50 - 100$ GeV, $\sim 80\%$ of the charged track energy is contained in a cone of phase space radius $R = \sqrt{\delta\eta^2 + \delta\phi^2} \sim 0.2$ [40], while an area of this size in a central Pb+Pb collision at 5.5 TeV may contain ~ 75 GeV [41] of uncorrelated energy from soft particle production. Simulation studies of jet reconstruction using unmodified fragmentation show that relatively small cone radii




Figure 5. Hard process rates at the LHC [39,45]. Solid and dashed lines for inclusive jets indicate rates vs E_T integrated over phase space cone of radius $R=0.4$ and 0.7 .

$R \sim 0.3-0.4$ provide optimum energy resolution [42], though the models of both jet signal (PYTHIA) and background (HIJING) in these studies do not capture the physics of jet softening and broadening due to medium effects. Full understanding of jet reconstruction capabilities and optimization of jet observables at the LHC must await data.

A recent calculation incorporating medium effects in the MLLA parton shower framework [43] predicts an enhancement in multiplicity due to jet quenching up to hadron $p_T \sim 5-6$ GeV/c for jets with $E_T \sim 100-200$ GeV. This excess should be measurable above background, enabling a detailed characterization of the modification of the fragmentation function. Measurement of high energy b-tagged and c-tagged jets can be contrasted to light hadron-led jets, which arise dominantly from gluons, to exploit the different color charge coupling of gluons and quarks to the medium (factor $9/4$). A broad range of multi-hadron correlations will be accessible which interpolate between leading hadron studies and full jet reconstruction.

The golden channel for jet quenching is the coincidence measurement of a jet recoiling from a gauge boson (γ or Z). The boson does not interact with the medium and therefore provides a clean calibration of the momentum transfer in the interaction [44], enabling measurement of the true fragmentation of the recoiling jet. While this strategy remains

attractive, the measurements are challenging. Fig. 5 shows that γ +jet rate is statistically robust in Pb+Pb only for $p_T < 40$ GeV/c. The γ/π^0 ratio (the key experimental parameter) exceeds 10% only for $p_T > 50$ GeV/c in p+p, though π^0 suppression may lower that bound to 20 GeV/c in central Pb+Pb [45]. Most significantly, QCD fragmentation photons may dominate the prompt photon yield up to 50 GeV or higher [46], complicating the interpretation of the photon as a non-interacting messenger from the hard vertex. The Z+jet channel is background-free but suffers from small cross section (see Fig. 5) and will be statistically marginal at nominal Pb+Pb luminosity.

6. Summary

The SPS and RHIC data presented at this conference significantly extend previous jet quenching studies, in some cases providing qualitatively new insights. The data are in broad agreement with expectations based on the dominant paradigm of radiative energy loss, but experimental tests of this correspondence are not yet definitive.

A characteristic feature of radiative energy loss is the quadratic path length dependence that arises from coherence effects. The new Cu+Cu data, combined with existing Au+Au measurements, will provide the most precise tests of this scaling. First hadron suppression measurements in Cu+Cu are in rough agreement with radiative energy loss models, but theoretical uncertainties now dominate this comparison and further progress will come from the theory side.

New di-hadron correlation studies reveal a well-defined recoil peak at high p_T , enabling the first differential measurements of jet suppression in the medium. Many existing calculations miss essential features of these data, perhaps due to approximations in the treatment of nuclear geometry or energy loss. Comparison to the most complete calculation available [36] gives a new constraint $\hat{q} \sim 5 - 7$ GeV²/fm, though the uncertainty on this number is difficult to estimate. Heavy flavor suppression, measured via non-photonic electrons, appears to be larger than expected from radiative energy loss alone [47], raising the question of significant collisional (elastic) energy loss.

The experimental study of jet quenching has reached a new level of detail and precision, and interpretation of the striking effects that have been observed is currently limited by theoretical uncertainties. Heavy ion collisions at the LHC will open up a huge new kinematic regime for jet quenching studies, with qualitatively new observables available. While the LHC regime may provide better grounds for quantitative theoretical predictions, a complete picture of jet interactions with dense QCD matter must describe all of the phenomena we observe, both at RHIC and at the LHC.

REFERENCES

1. J. Adams *et al.* [STAR Collaboration], Phys. Rev. Lett. 91 (2003) 172302.
2. S. S. Adler *et al.* [PHENIX Collaboration], Phys. Rev. Lett. 91 (2003) 072301.
3. C. Adler *et al.* [STAR Collaboration], Phys. Rev. Lett. 90 (2003) 082302.
4. X. N. Wang, M. Gyulassy and M. Plumer, Phys. Rev. D51 (1995) 3436.
5. R. Baier, Y. L. Dokshitzer, S. Peigne and D. Schiff, Phys. Lett. B345 (1995) 277.
6. M. Gyulassy, P. Levai and I. Vitev, Phys. Rev. Lett. 85 (2000) 5535.
7. X.-N. Wang and X. F. Guo, Nucl. Phys. A 696 (2001) 788.

8. R. Baier, D. Schiff and B. G. Zakharov, *Ann. Rev. Nucl. Part. Sci.* 50 (2000) 37.
9. C. Salgado and U. Wiedemann, *Phys. Rev. D* 68 (2003) 014008.
10. M. Mustafa and M. Thoma, *Acta Phys. Hung.* A22 (2005) 93.
11. M. Aggarwal *et al.* [WA98 Collaboration], *Eur. Phys. J.* C23 (2002) 225.
12. I. Vitev and M. Gyulassy, *Phys. Rev. Lett.* 89 (2002) 252301.
13. X.-N. Wang, *Phys. Rev. C* 61 89 (2000) 064910.
14. D. d'Enterria, *nucl-ex/0504001*.
15. A. Dainese [NA57 Collaboration], these proceedings.
16. A. Laszo and T. Schuster [NA49 Collaboration], these proceedings.
17. O. Barannikova [STAR collaboration], these proceedings.
18. S. S. Adler *et al.* [PHENIX Collaboration], *Phys. Rev. C* 69 (2004) 034909.
19. P. Sorensen, these proceedings.
20. N.N. Ajitanand [PHENIX Collaboration], these proceedings.
21. R. C. Hwa, these proceedings.
22. K. Eskola *et al.*, *Nucl. Phys.* A747 (2005) 511.
23. C. Salgado, these proceedings; *hep-ph/0510062*.
24. M. Shimomura, these proceedings; *nucl-ex/0510023*.
25. J. Dunlop [STAR Collaboration], these proceedings.
26. A. Dainese, C. Loizides and G. Paic, *Eur. Phys. J.* C38 (2005) 461.
27. R. Baier, *Nucl. Phys.* A715 (2003) 209c.
28. J. Adams *et al.* [STAR Collaboration], *Phys. Rev. Lett.* 95 (2005) 152301.
29. J. G. Ulery [STAR Collaboration], these proceedings.
30. N. Grau [PHENIX Collaboration], these proceedings.
31. D. Magestro [STAR Collaboration], these proceedings; *nucl-ex/0510002*.
32. X.-N. Wang, *Phys. Lett.* B595 (2004) 165.
33. A. Majumder, E. Wang and X.-N. Wang, *nucl-th/0412061*.
34. R. Baier *et al.*, *Nucl. Phys.* B484 (1997) 265.
35. I. Vitev, *hep-ph/0501225*.
36. C. Loizides *et al.*, these proceedings.
37. B. Muller and K. Rajagopal, *hep-ph/0502174*.
38. H.-A. Gustafsson, these proceedings.
39. C. Loizides, *nucl-ex/0501017*; A. Acardi *et al.*, *hep-ph/0310274*; N. Carrer and A. Dainese, *hep-ph/0311225*; N. Armesto, private communication.
40. T. Affolder *et al.* [CDF Collaboration], *Phys. Rev. D* 65 (2002) 092002.
41. K. Eskola, *et al.*, *hep-ph/0506049*.
42. S.-L. Blyth [ALICE-USA collaboration], *J. Phys.* G30 (2004) S1155.
43. N. Borghini and U. Wiedemann, these proceedings; *hep-ph/0506218*.
44. X.-N. Wang and Z. Huang, *Phys. Rev. C* 55 (1997) 3047.
45. F. Arleo *et al.*, *hep-ph/0311131*.
46. F. Arleo *et al.*, *JHEP* 11 (2004) 009.
47. J. Bielcik [STAR Collaboration]; S.A. Butsyk [PHENIX collaboration]; N. Armesto; and M. R. Djordjevic; these proceedings