

# High- $p_T$ $\pi^0, \eta$ , identified and inclusive charged hadron spectra from PHENIX

M. Shimomura<sup>a</sup> (for the PHENIX\* Collaboration)

<sup>a</sup>Graduate School of Pure and Applied Sciences, Univ. of Tsukuba, Tenno-dai 1-1-1, Tsukuba, Ibaraki, Japan

PHENIX has extended the measurement of the  $\pi^0, \eta$ , identified and inclusive charged hadrons up to 20 GeV/c, and extended the measurement to the Cu+Cu collision system. A strong suppression is observed for both  $\pi^0$  and charged hadron yields in central Au+Au and Cu+Cu collisions. Both in the Au+Au and Cu+Cu systems  $R_{AA}$  becomes independent of  $p_T$  above 5 GeV/c. Its centrality dependence is compared to two models in order to test for universal  $N_{part}$  scaling that is independent of system; results are inconclusive. The results are compatible with energy loss predictions. In addition, the ratio of  $\eta$  to  $\pi^0$  approaches, within uncertainties, a constant value of  $0.4 \sim 0.5$  at high  $p_T$  in  $p+p, d+Au$ , and Au+Au, while the ratio of  $K_s$  to  $\pi^0$  is also consistent with a constant value at high  $p_T$  in  $d+Au$  and  $p+p$ . These results are compatible with normal jet fragmentation.

## 1. PHYSICS MOTIVATION

We have previously observed that  $\pi^0, \eta$  and charged hadron yields are significantly suppressed especially for the high  $p_T$  region ( $p_T \geq 4 \sim 5$  GeV/c) in Au+Au collision at 200 GeV compared with  $p+p$  collisions [1–4]. Since there is no suppression in  $d+Au$  collisions at high  $p_T$  [3], it is understood that the suppression occurs due to some final state interaction in the collision such as gluon radiation in the hot dense matter. Another evidence for the suppression being a final state effect comes from the non-suppression of the direct photon yield in Au+Au collisions [5]. In order to better understand the character of the suppression, measurements made with different size systems ( $p+p/d+Au/Cu+Cu/Au+Au$ ) have been compared. Also, the large Au+Au dataset collected in Run-4 (2004) allowed us to extend the  $p_T$  reach to 20 GeV/c.

## 2. DATA ANALYSIS, $\pi^0, K_s$ , CHARGED HADRON AND $\eta$

We made new measurements of the following spectra.

- $\pi^0$  spectra with extended  $p_T$  range in Au+Au at 200GeV
- $\pi^0$  and charged hadron spectra in Cu+Cu at 200GeV
- $K_s$  spectra in  $d+Au$  and  $p+p$  at 200GeV [6]

---

\*For the full list of PHENIX authors and acknowledgements, see Appendix 'Collaborations' of this volume.

- Newly finalized  $\eta$  spectra in  $p+p$ ,  $d+Au$ , and  $Au+Au$  [7]

The PHENIX experiment consists of four spectrometer arms (two central arms and two muon arms) and a set of global detectors. Each central arm covers the pseudorapidity range  $|\eta| \leq 0.35$  and 90 degrees in azimuth. Charged particles are tracked by a drift chamber (DC) and pad chambers (PC) in each central arms. The electromagnetic calorimeters (EMCal) are used to measure  $\gamma$  energy deposit and construct the invariant masses of  $\pi^0(\rightarrow 2\gamma)$ ,  $K_s(\rightarrow 2\pi^0 \rightarrow 4\gamma)$  and  $\eta(\rightarrow 2\gamma)$  [4][8].

### 3. THE NUCLEAR MODIFICATION FACTOR $R_{AA}$

#### 3.1. Charged Hadron, $\pi^0$ and $\eta$

Fig. 1 shows the comparison of  $R_{AA}$  for  $\pi^0$  and charged hadrons in 0-10 % most central  $Au+Au$  and  $Cu+Cu$  collisions as function of  $p_T$ . Both  $\pi^0$  and charged hadron are strongly suppressed in both  $Au+Au$  and  $Cu+Cu$  collision. The difference between  $\pi^0$  and charged hadron for  $p_T \leq 5$  GeV/c comes from the proton contribution. For more central collisions, the suppression is getting stronger and the difference between  $\pi^0$  and charged hadron is getting larger. Above 5 GeV/c,  $\pi^0 R_{AA}$  becomes flat out to 20 GeV/c. These results are consistent with the model predicting parton energy loss in the medium [9] and the model predicting shadowing, Cronin effect, and parton energy loss in the medium [10]. In addition,  $\eta$  is also suppressed in central  $Au+Au$  collisions and the suppression pattern is similar to  $\pi^0$  [7].




Figure 1. The comparison of  $R_{AA}$  for  $\pi^0$  and charged hadron at 0-10 % centrality bin in  $Au+Au$  (left) and  $Cu+Cu$  (right) collisions as a function of  $p_T$  with theoretical prediction (purple [9] and red [10] lines). Error bars are statistical, boxes are the systematic error.

#### 3.2. Comparison between $Au+Au$ and $Cu+Cu$

In Fig 2(a), the comparison of  $R_{AA}$  in  $Au+Au$  to that in  $Cu+Cu$  with similar  $N_{part}$  is shown. The suppression is similar for similar  $N_{part}$  at mid-centrality. Although a universal  $N_{part}$  scaling (that holds at least approximately) has been suggested [11], due to the magnitude of our uncertainties it is still unclear whether a single scaling curve can exactly describe the suppression ( $R_{AA}$ ) in both systems simultaneously. In Fig. 2(b), we show the integrated  $R_{AA}$  of  $\pi^0$  at  $7.0 \leq p_T \leq 20.0$  GeV/c with two different theoretical curves [10][12] as a function of  $N_{part}$ . Both models are consistent with the data from mid-central to central collisions.




Figure 2. (a) The comparison between  $\pi^0$   $R_{AA}$  in Au+Au and Cu+Cu at similar  $N_{part}$  ( $\sim 74$ ). (b) The integrated  $R_{AA}$  at  $7 \leq p_T \leq 20$  GeV/c with theoretical curves [10][12] as a function of  $N_{part}$ .

#### 4. PARTICLE RATIOS

The ratio of  $\eta$  to  $\pi^0$  is  $\sim 0.4 - 0.5$  in all systems and for all centralities as shown in Fig. 3 [7]. Also, the ratio of  $K_s$  to  $\pi^0$  at  $p+p$  and  $d+Au$  becomes flat at high  $p_T$  as shown in Fig. 4 [6]. Therefore, the mesons are affected by the medium in the same way in different collision systems. These results are consistent with jet fragmentation at high  $p_T$ .




Figure 3. The ratio of  $\eta$  to  $\pi^0$  in Au+Au (left),  $d+Au$  (right) and  $p+p$  (right) as a function of  $p_T$  at  $\sqrt{s} = 200$  GeV. Error bars are statistical error, boxes are the systematic error.




Figure 4. Closed symbols are the ratio of  $K_s$  to  $\pi^0$  in  $p+p$  and  $d+Au$  as a function of  $p_T$ . Open symbols show the ratio of  $K^\pm$  to  $\pi^\pm$  as reference [13]. The error bars are statistical error, and the boxes are systematic error.

## 5. SUMMARY

We have studied  $\pi^0, \eta, K_s$  and charged hadron spectra in Au+Au, Cu+Cu,  $d+Au$  and  $p+p$  at high  $p_T$ . For  $\pi^0$  and charged hadron, we observed the suppressions in both Cu+Cu and Au+Au collisions compared with  $p+p$  collisions, and no suppression is observed in  $d+Au$  collisions. The  $R_{AA}$  comparison between Au+Au and Cu+Cu indicates that the suppression is almost the same for similar  $N_{part}$ . A universal  $N_{part}$  scaling of  $R_{AA}$ , independent of system, describes the data in an approximate sense, but it cannot be confirmed exactly due to experimental uncertainties. In addition, the high  $p_T$   $\pi^0$  suppression is flat out to 20 GeV/c and its magnitude is quantitatively consistent with parton energy loss model calculations.  $\eta$  has a similar suppression pattern as  $\pi^0$  does. The ratio of  $\pi^0$  to  $\eta$  is independent of centralities and system size. Similarly, the  $K_s$  to  $\pi^0$  ratio is constant within uncertainties at high  $p_T$  for both  $p+p$  and  $d+Au$ . These particle ratios are consistent with normal jet fragmentation.

## REFERENCES

1. S. S Adler *et al.*, PHENIX Collaboration, Phys. Rev. Lett. 91, 241803 (2003).
2. S. S Adler *et al.*, PHENIX Collaboration, Phys. Rev. Lett. 91, 072301 (2003).
3. S. S Adler *et al.*, PHENIX Collaboration, Phys. Rev. Lett. 91, 072303 (2003).
4. S. S Adler *et al.*, PHENIX Collaboration, Phys. Rev. C 69, 034910 (2004).
5. S. S Adler *et al.*, PHENIX Collaboration, Phys. Rev. Lett. 94, 232301 (2005).
6. V. Ryabov for the PHENIX Collaboration, (in this Proceedings)
7. S. S Adler *et al.*, PHENIX Collaboration, Update of "High  $p_T$   $\eta$  meson production in  $p+p$ ,  $d+Au$  and Au+Au collisions at  $\sqrt{s_{NN}} = 200$  GeV", to be submitted to PLB
8. K. Adcox *et al.*, Nucl. Instrum. Methods A499, 469 (2003).
9. X. Wang, Phys. Lett. B 595, 165-170 (2004).
10. I. Vitev, Phys. Rev. Lett. 89, 252301 (2002).
11. A. Drees, H. Feng, J. Jia, Phys. Rev. C 71, 034909 (2005).
12. V. S. Pantuev, hep-ph/0506095 (2005).
13. S. S Adler *et al.*, PHENIX Collaboration, Update of "Identified charged hadron production in  $\sqrt{s} = 200$  GeV  $p+p$  collisions", to be submitted to PRD